



# Intake of *Lactobacillus reuteri* Improves Incretin and Insulin Secretion in Glucose-Tolerant Humans: A Proof of Concept

Diabetes Care 2015;38:1827–1834 | DOI: 10.2337/dc14-2690

Marie-Christine Simon,<sup>1,2</sup>  
 Klaus Strassburger,<sup>2,3</sup> Bettina Nowotny,<sup>1,2</sup>  
 Hubert Kolb,<sup>4</sup> Peter Nowotny,<sup>1,2</sup>  
 Volker Burkart,<sup>1,2</sup> Fariba Zivehe,<sup>1,2</sup> Jong-  
 Hee Hwang,<sup>1,2</sup> Peter Stehle,<sup>5</sup>  
 Giovanni Pacini,<sup>6</sup> Bolette Hartmann,<sup>7</sup>  
 Jens J. Holst,<sup>7</sup> Colin MacKenzie,<sup>8</sup>  
 Laure B. Bindels,<sup>9</sup> Ines Martinez,<sup>9</sup>  
 Jens Walter,<sup>9,10,11</sup> Birgit Henrich,<sup>8</sup>  
 Nanette C. Schloot,<sup>1,2</sup> and  
 Michael Roden<sup>1,2,12</sup>

## OBJECTIVE

Ingestion of probiotics can modify gut microbiota and alter insulin resistance and diabetes development in rodents. We hypothesized that daily intake of *Lactobacillus reuteri* increases insulin sensitivity by changing cytokine release and insulin secretion via modulation of the release of glucagon-like peptides (GLP)-1 and -2.

## RESEARCH DESIGN AND METHODS

A prospective, double-blind, randomized trial was performed in 21 glucose-tolerant humans (11 lean: age  $49 \pm 7$  years, BMI  $23.6 \pm 1.7$  kg/m<sup>2</sup>; 10 obese: age  $51 \pm 7$  years, BMI  $35.5 \pm 4.9$  kg/m<sup>2</sup>). Participants ingested  $10^{10}$  b.i.d. *L. reuteri* SD5865 or placebo over 4 weeks. Oral glucose tolerance and isoglycemic glucose infusion tests were used to assess incretin effect and GLP-1 and GLP-2 secretion, and euglycemic-hyperinsulinemic clamps with [6,6-<sup>2</sup>H<sub>2</sub>]glucose were used to measure peripheral insulin sensitivity and endogenous glucose production. Muscle and hepatic lipid contents were assessed by <sup>1</sup>H-magnetic resonance spectroscopy, and immune status, cytokines, and endotoxin were measured with specific assays.

## RESULTS

In glucose-tolerant volunteers, daily administration of *L. reuteri* SD5865 increased glucose-stimulated GLP-1 and GLP-2 release by 76% ( $P < 0.01$ ) and 43% ( $P < 0.01$ ), respectively, compared with placebo, along with 49% higher insulin ( $P < 0.05$ ) and 55% higher C-peptide secretion ( $P < 0.05$ ). However, the intervention did not alter peripheral and hepatic insulin sensitivity, body mass, ectopic fat content, or circulating cytokines.

## CONCLUSIONS

Enrichment of gut microbiota with *L. reuteri* increases insulin secretion, possibly due to augmented incretin release, but does not directly affect insulin sensitivity or body fat distribution. This suggests that oral ingestion of one specific strain may serve as a novel therapeutic approach to improve glucose-dependent insulin release.

Type 2 diabetes results from decreased insulin sensitivity and inadequate insulin secretion, which associate with diminished incretin response and subclinical chronic inflammation and subsequent impaired glucose tolerance (1–4). These pathogenic factors, frequently accompanied by hypercaloric high-fat low-fiber diets, may be associated with alterations in gut microbiota, which also occur in obesity (5) and type 2 diabetes (6).

<sup>1</sup>Institute for Clinical Diabetology, German Diabetes Center, Leibniz Center for Diabetes Research, Heinrich-Heine University, Düsseldorf, Germany

<sup>2</sup>German Center for Diabetes Research, Düsseldorf, Germany

<sup>3</sup>Institute for Biometry and Epidemiology, German Diabetes Center, Leibniz Center for Diabetes Research, Heinrich-Heine University, Düsseldorf, Germany

<sup>4</sup>West-German Centre of Diabetes and Health, Verbund Katholischer Kliniken Düsseldorf, Düsseldorf, Germany

<sup>5</sup>Department of Nutrition and Food Science, Nutritional Physiology, Rheinische Friedrich-Wilhelms-University, Bonn, Germany

<sup>6</sup>Metabolic Unit, Institute of Biomedical Engineering, National Research Council, Padova, Italy

<sup>7</sup>Novo Nordisk Foundation Center for Basic Metabolic Research, Department for Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark

<sup>8</sup>Institute for Medical Microbiology and Hospital Hygiene, Heinrich-Heine University, Düsseldorf, Germany

<sup>9</sup>Department of Food Science and Technology, University of Nebraska-Lincoln, Lincoln, NE

<sup>10</sup>Department of Agricultural, Nutritional and Food Science, University of Alberta, AB, Canada

<sup>11</sup>Department of Biological Sciences, University of Alberta, Edmonton, Canada

<sup>12</sup>Department of Endocrinology and Diabetology, Medical Faculty, Heinrich-Heine University, Düsseldorf, Germany

Corresponding authors: Michael Roden, michael.roden@ddz.uni-duesseldorf.de, and Nanette C. Schloot, schloot@ddz.uni-duesseldorf.de.

Received 12 November 2014 and accepted 1 May 2015.

Clinical trial reg. no. NCT01250106, clinicaltrials.gov.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc14-2690/-/DC1>.

A slide set summarizing this article is available online.

N.C.S. and M.R. contributed equally.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

See accompanying article, p. 1817.

In animal models, modulation of intestinal microbiota may contribute to regulation of insulin resistance and postprandial glucose response (7). Mice treated with prebiotics exhibited reduced gut permeability and increased plasma glucagon-like peptide (GLP)-2 concentrations, which is cosecreted with GLP-1 and exerts an intestinotrophic effect. The improved gut barrier function further correlated with lower portal plasma lipopolysaccharide (LPS) (endotoxin) concentrations and lower systemic and hepatic inflammation (8).

Previous studies in humans and animals reported conflicting results on the effects of different *Lactobacillus* strains on body weight (9). In humans, ingestion of prebiotics has been shown to influence gut microbiota composition (10), perhaps alter intestinal permeability, and increase plasma GLP-1 concentrations and decrease postprandial glucose concentrations (11). Another recent study investigated the effects of allogenic fecal transplantation on glucose metabolism. Infusion of fecal preparations from lean donors transiently reduced insulin resistance in obese individuals with metabolic syndrome (12). Ingestion of *Lactobacillus acidophilus* NCFM for 4 weeks maintained insulin sensitivity without effect on the systemic inflammatory response in a metabolically heterogeneous group (13). Consumption of a yogurt containing *L. acidophilus* La5 and *Bifidobacterium lactis* Bb12 over 6 weeks improved oxidative and glucose metabolism in patients with type 2 diabetes (14). While the latter studies suggested a beneficial metabolic action on the host by *Lactobacilli*, even without modulating the microbiome (15), two double-blinded studies in obese adolescents failed to detect any beneficial effects of probiotics on the metabolic syndrome (16,17). At present, a randomized controlled intervention trial including analyses of tissue-specific effects and the possible underlying mechanisms is missing in metabolically well-phenotyped humans. Thus, this proof-of-principle study aimed at detailed examination of the effects of *Lactobacilli* on insulin secretion and action in humans. For this initial approach, healthy lean and obese individuals were recruited to test the effects without interference from glucose-lowering medication and variable degree of gut microbial dysbiosis (18,19).

We hypothesized that *L. reuteri* supplementation improves insulin sensitivity and increases insulin secretion by augmenting incretin release. Thus, we investigated the effect of ingested *L. reuteri* on incretin, tissue-specific insulin sensitivity, and  $\beta$ -cell function using isoglycemic and euglycemic clamps and measured fat distribution as well as markers of systemic inflammation in healthy lean and obese individuals. *L. reuteri* was selected for the intervention because of its well-studied safety and its strong probiotic activity compared with 46 other strains of *Lactobacillus* spp. (15,20).

## RESEARCH DESIGN AND METHODS

### Study Design

We performed a double-blind, 1:1 randomized, prospective, longitudinal pilot trial over 8 weeks. After a run-in phase, volunteers received BIO-tract controlled-release caplets with *L. reuteri* SD5865 at a dose of  $2 \times 10^{10}$  viable cells, encapsulated by Nutraceutix (Nutraceutix, Inc., Redmond, WA) and packaged identical to the placebo controls. Study participants started with a run-in phase of 4 weeks after the baseline visit during which they received placebo only. Then, they received capsules containing placebo or  $10^{10}$  cells of *L. reuteri* to be ingested b.i.d. for 4 consecutive weeks.

Anthropometric measurements, oral glucose tolerance tests (OGTT), clamp studies, liver and muscle fat measurements, and collection of fecal samples were performed at the end of the run-in phase and at end of the trial. At each visit, participants underwent interviews regarding adverse effects, symptoms, or changes in quality of life. Compliance was assessed by counting capsules and by screening for *L. reuteri* in gut microbiota using quantitative real-time PCR. The study was performed according to the Declaration of Helsinki, approved by the local ethics committee, and registered at ClinicalTrials.gov. All participants provided written informed consent.

### Participants

Lean (BMI 19–25 kg/m<sup>2</sup>) and obese (30–45 kg/m<sup>2</sup>) participants matched for age and sex were recruited at the German Diabetes Center. Inclusion criteria were age 40–65 years, nonsmoking, absence of gastrointestinal disease, and willingness to abstain from intake of other

probiotic food and fermented milk products over the study period but otherwise stick to previous eating habits. Exclusion criteria were pregnancy, cancer, chronic diseases, treatment with antibiotics within 3 months prior to study inclusion and throughout study, competitive athletics, impaired glucose tolerance, and impaired fasting glucose.

### Anthropometric Measurements

Body weight, height, waist circumference, and blood pressure were measured at each visit. Bioelectrical impedance analysis (Nutriguard-S; Data Input GmbH, Darmstadt, Germany) was performed in duplicates, and lean body and fat mass were calculated (21).

### Gastric Emptying

Each subject underwent a [<sup>13</sup>C]octanoic acid breath test to measure gastric emptying (22). In brief, the appearance of <sup>13</sup>CO<sub>2</sub> in breath after oral administration of <sup>13</sup>C-octanoic acid applied with egg yolk was determined. Data were used to assess disturbed gastric emptying, as there is a reciprocal interplay of gut hormones and gastric motility.

### OGTT and Isoglycemic Clamp Test

OGTT was performed after an overnight fast with 75 g glucose and 300 mL water (Dextrose O.G-T.; Roche Diagnostics, GmbH, Mannheim, Germany).  $\beta$ -Cell function was assessed from the insulinogenic index (23) and insulin sensitivity from the oral glucose insulin sensitivity index (OGIS) (24), which provides a measurement of insulin-mediated glucose clearance. The interplay between insulin sensitivity and secretion, which describes the  $\beta$ -cell adaptive response to changes of insulin resistance (25), was determined by the products  $OGIS \times AUC_{CP}$  and  $OGIS \times AUC_{INS}$ , where the AUCs are the area under the concentration curves of C-peptide and insulin, respectively. Adaptation and disposition index were originally developed for the intravenous glucose test but proved useful determinants of  $\beta$ -cell function also for OGTT (26). Further, the glucose AUC was used to measure differences in glucose tolerance upon intervention.

For measurement of the incretin effect, isoglycemic intravenous glucose infusion (20% in sterile water; Braun, Melsungen, Germany) was performed, after an overnight fast, over 3 h aiming at mimicking the blood glucose level measured during OGTT (27).

### Euglycemic-Hyperinsulinemic Clamp Test

The clamp test was combined with isotopic dilution ([6,6-<sup>2</sup>H<sub>2</sub>]glucose) to assess both whole-body insulin sensitivity and hepatic insulin sensitivity. Constant insulin infusion (40 mU/m<sup>2</sup> body surface area i.v.; Insuman Rapid, Sanofi, Frankfurt, Germany) was given over 2 h, and blood glucose was adjusted to 80 mg/dL by variable 20% glucose infusion. Glucose infusion rate during the last 30 min during the clamp were used to calculate insulin sensitivity. Rates of endogenous glucose production (EGP) were determined from the tracer infusion rate of [6,6-<sup>2</sup>H<sub>2</sub>]glucose and its enrichment to hydrogens bound to carbon 6 divided by the mean percent enrichment of plasma [6,6-<sup>2</sup>H<sub>2</sub>]glucose. Steady-state equations were considered appropriate for calculation of basal EGP and insulin-suppressed EGP during the last 30 min of the clamp (28).

### Indirect Calorimetry

For determination of in vivo whole-body energy expenditure and substrate oxidation from measuring oxygen consumption and carbon dioxide production, indirect calorimetry was performed, using Vmax Encore 29n (CareFusion, Höchberg, Germany) as previously described (28).

### Determination of Ectopic Fat by

**<sup>1</sup>H-Magnetic Resonance Spectroscopy** Measurements were performed on a 3.0 Tesla magnet (X series; Achieva, Philips, Best, the Netherlands). Ectopic fat content from <sup>1</sup>H-magnetic resonance spectroscopy was quantified relative to internal water content and calculated as previously described (29). Data were analyzed using jMRUI software. Ectopic lipid content was assessed in eight lean (4 *L. reuteri* and 4 placebo) and seven obese (4 *L. reuteri* and 3 placebo) volunteers.

### Analysis of Fecal Microbiota Composition and Content

For exact quantification of specific taxa, we performed quantitative real-time PCRs to detect total bacterial load (eubacteria), enterobacteria content, *L. spp.*, and *L. reuteri*. In brief, total genomic DNA of the fecal samples was prepared within 24 h after collection and stored at -20°C until further analysis. DNA extraction was performed by using a BioRobot EZ1 machine

(Qiagen, Hilden, Germany). Primers and probes used for qPCR were retrieved from the literature ("eubacteria" [30], "*Lac. spp.*" [31], "*L. reuteri*" [32]) or designed by Primer Express software (DNASTAR, Madison, WI) for enterobacteria content. Next-generation sequencing for analysis of fecal content was performed in cooperation with the University of Minnesota Genomics Center as described in detail in Supplementary Material 1.

### Laboratory Measurements

Blood cell count, HbA<sub>1c</sub>, hs-CRP, and blood glucose were measured by routine methods as previously described (28). Serum C-peptide, insulin, and plasma glucagon were measured by RIA (Millipore, St. Charles, MO). Cytokines and chemokines were assayed by the Flurokine MAP multiplex kit method (R&D Systems, Minneapolis, MN) and Luminex technology (Bio-Rad Bioplex, Nazareth, Belgium) (33). LPS was quantified using a chromogenic limulus amoebocyte lysate QCL-1000 assay (Lonza, Ltd., Verviers, Belgium) according to the manufacturer's instructions. For validation of the assay, samples were spiked with known amounts of LPS and yielded a mean recovery rate of 90.4% (34). Production of reactive oxidative species was estimated from serum thiobarbituric acid-reactive substances (35). For incretin measurements, blood was collected into dipeptidyl peptidase-4 inhibitor-containing tubes (BD P700; BD Biosciences, Franklin Lakes, NJ). GLP-1 (7-36)amide and GLP-1 (9-36)amide and only fully processed GLP-2 of intestinal origin were measured as previously described (36,37). After deproteinization, determination of atom percent enrichment of <sup>2</sup>H was performed by gas chromatography-mass spectrometry as previously described (28).

### Calculations and Statistics

Data are presented as mean ± SD or medians and interquartile range as appropriate. Variables with skewed distribution were log transformed before further analysis. To test differences between treatment arms taking the lean-obese status into account, we used two-way ANOVA. *P* values from two-sided tests ≤5% were considered to indicate significant differences. SAS for Windows version 9.2 (SAS Institute, Cary, NC) was used (Supplementary Material 2). For the individual hormones and metabolites,

the total AUC was calculated using the trapezoidal rule. The incretin effect was assessed from comparing the total AUC values for insulin, C-peptide, glucagon, GLP-1, and GLP-2 during OGTT and with those obtained during the isoglycemic test. The ΔAUC was defined as the difference between the AUCs during OGTT and the isoglycemic test (Supplementary Material 2). Because of skewed distribution, the ΔAUCs were log transformed (natural logarithm). Differences of these ΔAUC values obtained before and after intervention were statistically analyzed.

## RESULTS

### Study Population

Thirty-nine persons were screened from October 2011 until August 2012. Thirteen persons did not pass the inclusion criteria. Of the remaining 21 participants, 5 lean received placebo and 6 lean received *L. reuteri*, whereas 5 obese received placebo and 5 obese received *L. reuteri* (Supplementary Fig. 1). Treatment adherence was 100% in all study groups (Table 1). None of the participants reported any adverse effects including gastrointestinal disorders.

### Gut Microbiota Analyses

Fecal samples were obtained from all 21 participants before intervention and from all but 1 subject at the end of intervention. *L. reuteri* was not detectable before intervention in all but one lean subject of the placebo group. This subject was tested weakly positive (0.249 DNA copies) before intervention but lower after the intervention (0.041 DNA copies). All participants of the group treated with *L. reuteri* tested positive after intervention, while those of the placebo group were negative (*P* < 0.0001). In all groups, both treatments did not alter the total bacterial load, the enterobacteria, or the levels of total *Lactobacilli* (Supplementary Fig. 2).

Analysis of fecal microbiota composition by next-generation sequencing revealed that *L. reuteri* administration did not affect the overall composition of the fecal microbiota. No significant differences were observed when comparing temporal changes in relative abundance of bacterial taxa or operational taxonomic units between placebo-treated and *L. reuteri*-treated participants either when all participants were considered or when they were divided into lean and

**Table 1—Baseline characteristics of study participants**

	Combined	Lean	Obese
N (female/male)	11/10	6/5	5/5
Age (years)	50 ± 7	49 ± 7	51 ± 7
BMI (kg/m <sup>2</sup> )	29.2 ± 7.0	<b>23.6 ± 1.7</b>	<b>35.5 ± 4.9*</b>
Waist circumference (cm)	99.1 ± 17.7	<b>83.3 ± 11.5</b>	<b>111.9 ± 9.0*</b>
Body fat (%)	34 ± 9	<b>28 ± 6</b>	<b>40 ± 9*</b>
Lean body mass (kg)	56 ± 12	<b>50.7 ± 11.1</b>	<b>63.5 ± 9.5*</b>
Fasting blood glucose (mg/dL)	78.3 ± 6.8	77 ± 9	79 ± 6
Fasting free fatty acids (μmol/L)	439 (370–631)	439 (246–522)	462 (386–678)
OGIS (mL/min/m <sup>2</sup> )	528 ± 92	<b>569 ± 94</b>	<b>482 ± 67*</b>
Adaptation index	0.808 (0.511–0.924)	0.615 (0.509–0.924)	0.830 (0.720–0.957)
Disposition index	3.09 (2.68–5.09)	2.76 (2.33–3.08)	4.77 (4.33–7.55)
Insulinogenic index	179 (123–396)	135 (94–409)	219 (147–395)
hs-CRP (mg/dL)	0.09 (0.05–0.29)	<b>0.05 (0.04–0.10)</b>	<b>0.30 (0.08–0.40)*</b>
Systolic blood pressure (mmHg)	138 ± 21	133 ± 23	143 ± 16
Diastolic blood pressure (mmHg)	85 ± 11	<b>77 ± 9</b>	<b>93 ± 6*</b>
Gastric emptying (t1/2)#	84 ± 34	90 ± 36	78 ± 32
Treatment adherence†	21/21	11/11	10/10

Mean ± SD and median (interquartile range) are given for normal and log-normal distributed data, respectively. Significant differences (\* $P < 0.05$ ) as determined by unpaired  $t$  test between lean and obese persons are marked in bold. #Gastric emptying as measured by the <sup>13</sup>C-octanoic acid breath test is given as half-life (t1/2) in minutes. †Compliance was confirmed by tablet counting.

obese subgroups.  $\alpha$ -Diversity analysis showed that the *L. reuteri* treatment did not modify the richness or evenness of the ecosystem. Principal coordinate analysis of  $\beta$ -diversity distance metrics did not show clustering due to the treatment (Supplementary Fig. 2). On an individual basis, these principal coordinate analyses suggest that, for most individuals, their fecal microbiota was relatively stable over time, as previously reported (38).

#### Anthropometry

The obese group had higher BMI ( $P < 0.001$ ), body fat and lean body mass, hs-CRP, and diastolic blood pressure but lower OGIS in line with insulin resistance (Table 1). Body weight and resting energy expenditure of lean and obese participants differed ( $P < 0.002$ ) but remained stable during the study and were not affected by the treatments (Supplementary Table 2) Waist-to-hip ratio and blood pressure also did not change (data not shown).

#### Ectopic Lipid Contents

Ingestion of *L. reuteri* did not affect hepatocellular lipids or intramyocellular lipids of m. soleus and m. tibialis anterior (Supplementary Table 2) in combined groups or in lean and obese subgroups. Hepatocellular lipids, but not intramyocellular lipids in both muscles, correlated with BMI before ( $r = 0.72$ ,  $P < 0.003$ ) and after ( $r = 0.63$ ,  $P < 0.014$ ).

#### Tissue-Specific Insulin Sensitivity and Substrate Oxidation

Whole-body insulin sensitivity was not affected by intervention. The *L. reuteri* and placebo groups showed no differences before or after intervention (Supplementary Fig. 3A). Steady-state conditions of glucose, insulin, C-peptide, and cortisol were comparable. Lean participants showed better whole-body insulin sensitivity compared with obese before ( $P < 0.05$ ) and after ( $P < 0.05$ ) intervention. In line with this, lipid oxidation rates during hyperinsulinemia were reduced by *L. reuteri* (94%) and placebo (102%) compared with baseline without differences between lean and obese. Oxidative glucose utilization and nonoxidative glucose utilization were increased during clamp but not affected by treatment (Supplementary Fig. 3B) and showed no differences in subgroup analysis for lean and obese.

*L. reuteri* did not affect EGP at baseline or during EGP suppression, being comparable between *L. reuteri* and placebo groups before and after intervention (Supplementary Fig. 3C) as well as in subgroup analysis of lean and obese.


#### Glucose Tolerance

Blood glucose levels during OGTT remained unchanged upon treatment (Supplementary Fig. 4A). Likewise, isoglycemic intravenous glucose infusion

(Supplementary Fig. 4B) and the total glucose infusion rate showed no differences in the intervention or placebo group. Glucose infusion rate of lean and obese persons differed after (median 13.0 [interquartile range 6.9–29.6] vs. 37.5 [22.0–47.9],  $P < 0.004$ ) but not before (23.2 [10.3–26.2] vs. 22.3 [19.3–51.3],  $P = 0.3073$ ) intervention, while subgroup-analysis of lean and obese participants showed no treatment effect.

#### Insulin Secretion and $\beta$ -Cell Function

After intervention  $\Delta$ AUC insulin and C-peptide were increased in the *L. reuteri* ( $P < 0.05$ ), but not the placebo group (Fig. 1A and B). The mean increase of the  $\Delta$ AUC insulin concentrations was 49% (46% for lean and 51% obese; significant only for combined group) and for C-peptide 55% (39% for lean and 74% obese; significant only for combined group). Increased concentrations were seen for insulin and C-peptide at 20 and 60 min of oral glucose stimulation in the *L. reuteri* group at the end of the trial ( $P < 0.05$ ) (Fig. 1A and B). Maximal response for insulin and C-peptide was increased at the end of the study in the intervention group ( $P < 0.016$ ,  $P < 0.018$ , respectively). Additionally, adaptation, disposition, and insulinogenic indices were increased in the *L. reuteri* group ( $P < 0.05$ ) upon treatment compared with placebo group (Supplementary Table 2).



**Figure 1**—Insulin, C-peptide, and glucagon. Insulin, C-peptide, and glucagon concentrations during OGTT before (open symbols) and after (closed symbols) intervention of *L. reuteri* and placebo groups, respectively. Shown are data from lean (triangles) and obese (circles) persons as mean ± SD. Inserts show combined group (lean and obese) ΔAUC before vs. after intervention of *L. reuteri* and placebo groups. Shown are boxes and whiskers. Error bars represent the minimum and maximum value. \* $P < 0.05$  vs. baseline for combined group.

Subgroup analysis showed differences upon treatment for the adaptation index ( $P < 0.022$ ) and the disposition index ( $P < 0.033$ ) only in the obese group.


**Incretin Response and Incretin Effect**  
Administration of oral glucose leads to increased levels of GLP-1 and GLP-2 within a maximum at 30 min for GLP-1 and 60 min for GLP-2 (Fig. 2A and B).

Maximal responses for GLP-1 and GLP-2 were increased at the end of the study in the intervention group for lean participants ( $P < 0.02$ ,  $P < 0.04$ , respectively) but not for obese participants.

Fasting plasma GLP-1 levels were different within the intervention group after treatment ( $P < 0.02$ , 0 min) and were by trend increased ( $P = 0.052$ , 0 min) compared with placebo. Glucose-stimulated GLP-1 levels were higher

after treatment ( $P < 0.05$ ) within the *L. reuteri* group (Fig. 2A) and compared with placebo ( $P < 0.05$ ). Similarly, glucose-stimulated GLP-2 levels tended to increase after treatment ( $P = 0.056$  and  $P = 0.065$ , respectively) within the *L. reuteri* group (Fig. 2B) and increased compared with placebo (20 min:  $P < 0.05$ , 60 min:  $P < 0.02$ ).

Glucose-stimulated AUC of GLP-1 was increased in the *L. reuteri* group after



**Figure 2**—GLP-1 and GLP-2 concentrations during OGTT. GLP-1 and GLP-2 concentrations during OGTT before (open symbols) and after (closed symbols) intervention. Shown are lean (triangles) and obese (circles) persons as mean  $\pm$  SEM. Inserts show combined group (lean and obese)  $\Delta$ AUC before vs. after intervention of *L. reuteri* and placebo groups. Shown are boxes and whiskers. Error bars represent the minimum and maximum value. \* $P < 0.05$ , \*\* $P < 0.01$  vs. baseline for combined group.

intervention ( $P < 0.003$ ) and enlarged compared with the placebo group (Fig. 2A). Similarly, glucose-stimulated AUC of GLP-2 was increased in the intervention group compared with placebo ( $P < 0.029$ ) (Fig. 2B).

$\Delta$ AUC of GLP-1 was increased at the end of the study compared with placebo ( $P < 0.004$ ) and tended to increase within the *L. reuteri* group ( $P = 0.087$ ) (Fig. 2A). *L. reuteri* increased  $\Delta$ AUC of GLP-2 ( $P < 0.02$ ) significantly different from placebo ( $P < 0.004$ ) (Fig. 2B).

Concentrations of GLP-1 and GLP-2 during OGTT showed a similar pattern and correlated positive intraindividually before (mean  $r = 0.69$ ;  $P < 0.0001$ ) and after (mean  $r = 0.73$ ;  $P < 0.0001$ ) intervention. Additionally, during OGTT GLP-1 and GLP-2 correlated intraindividually with insulin and C-peptide (all  $P < 0.001$ ) but not during the isoglycemic test (data not shown). Likewise, AUCs of GLP-1 and GLP-2 correlated before

( $r = 0.87$ ,  $P < 0.0001$ ) and after ( $r = 0.95$ ,  $P < 0.0001$ ) intervention.

#### Endotoxin, Cytokines, and Oxidative Stress

Fasting serum endotoxin levels were higher in obese compared with lean participants before intervention ( $P < 0.019$ ) but not afterward ( $P = 0.169$ ) (Supplementary Fig. 5). Serum endotoxin levels correlated with BMI before ( $r = 0.443$ ,  $P < 0.045$ ) but not after ( $r = 0.354$ ,  $P = 0.115$ ) intervention. Before and after intervention, obese participants had higher fasting hs-CRP ( $P < 0.004$  and  $P < 0.002$ ) as well as higher interleukin (IL)-1ra, MCP-1, and tumor necrosis factor- $\alpha$ . IL-8 and MIP-1 $\beta$  were not different between groups and interventions. Ingestion of *L. reuteri* did not affect the ratios between analyzed pro- and anti-inflammatory immune mediators, including the tumor necrosis factor- $\alpha$ -to-IL-1ra ratio. Measurements

of  $\gamma$ -interferon, IL-6, IL-1 $\beta$ , and IL-10 concentrations yielded  $>80\%$  of values below the detection limit and were therefore not further analyzed. Finally, serum thiobarbituric acid-reactive substances, as a measure of reactive oxygen species differed neither between groups nor between interventions (Supplementary Table 3).

#### CONCLUSIONS

This study demonstrates that ingestion of *L. reuteri* modulates the secretion of insulin, C-peptide, and proglucagon-derived gut peptides in humans, whereas insulin sensitivity and parameters influencing insulin sensitivity such as body mass and ectopic fat content, as well as systemic inflammation or oxidative stress, remain unchanged. Stool samples from persons ingesting *L. reuteri* were positive tested; placebo-treated persons stayed negative, while no differences were detected in other bacterial

members of the fecal microbiota. In line with our results in humans, Poutahidis et al. (15) showed that ingestion of *L. reuteri* has an effect on the host physiology of mice, altering immunological and metabolic aspects without changing the gut microbial ecology.

GLP-2 primarily targets intestinal mucosa promoting growth and intestinal integrity (8). Both GLP-1 and GLP-2 are processed from proglucagon with equal efficiency and secreted by the intestinal L cells in parallel and in equimolar amounts (37). Indeed, GLP-1 and GLP-2 measured in our study were positively correlated. GLP-1 and GLP-2 correlate intraindividually with insulin and C-peptide concentrations during OGTT but not during the isoglycemic test. The association of GLP-2 concentrations with those of insulin probably is indirect and explained by the cosecretion of GLP-2 with the hormone GLP-1. After treatment with *L. reuteri*, the mean increase of GLP-2  $\Delta$ AUC during OGTT was 43% compared with baseline and this was similar to the 76% increase seen for GLP-1  $\Delta$ AUC. An involvement of both peptides, the intestinotrophic GLP-2 and insulinotrophic GLP-1, in effects of a prebiotic treatment was found previously in animal models (8) and even in germ-free mice (39). Our study complemented these findings, showing that the probiotic strain *L. reuteri* increases the glucose-stimulated release of GLP-1 and GLP-2 in glucose-tolerant human participants.

The increased GLP-1 concentrations can serve to explain the increased glucose-stimulated insulin secretion in the intervention-group. The mean  $\Delta$ AUC insulin was increased by 49% (46% for lean, 51% obese). This was paralleled as expected by an increase of  $\Delta$ AUC C-peptide by 55% (39% for lean, 74% obese) and also accompanied by altered  $\beta$ -cell function, determined by insulinogenic adaptation and disposition index. At first sight, the increased insulin secretion observed in association with increased GLP-1 and  $\beta$ -cell function suggests that *L. reuteri* ingestion would offer a potential means to apply this regimen in patients with type 2 diabetes. However, as the investigated participants were healthy, whether the increased incretin and insulin concentrations obtained here can be extrapolated to patients with type 2 diabetes remains unresolved and is of physiological relevance.

There was no impact on hepatic and peripheral insulin sensitivity, glucagon concentrations, or hepatocellular and myocellular lipid content by *L. reuteri* administration. The only other study of the probiotic effect on insulin sensitivity reported that consumption of *L. acidophilus* NCFM for 4 weeks in a heterogeneous study population of 54 males with type 2 diabetes or with impaired or normal glucose tolerance revealed stable insulin sensitivity compared with the placebo group (13). There was considerable variability among study participants, and it was not reported whether those with an increase of insulin sensitivity were diabetic or normoglycemic. Only in the latter case would there be a difference from our observations in glucose-tolerant participants.

The relationship between insulin sensitivity, insulin level, and glucose tolerance is to some extent mediated by free fatty acid (4). However, there were no alterations of free fatty acid concentrations due to intervention (Supplementary Fig. 6), whereas *L. reuteri* changed incretin concentrations and insulin secretion.

At least one study suggested beneficial effects of probiotics on liver disease such as nonalcoholic fatty liver disease and alcoholic liver cirrhosis in humans (40). In our study, the hepatocellular and myocellular lipid content of all participants remained unchanged, which may again be due to the fact that we studied healthy persons.

Our study in healthy lean and obese persons with normal glucose tolerance did not show alterations of blood glucose or HbA<sub>1c</sub> levels. Interestingly, a recent prospective dietary analysis showed that greater intake of yogurt is associated with decreased risk of type 2 diabetes (41).

We also searched for additional metabolic effects of the ingestion of *L. reuteri* that involve improved incretin release. An increase of GLP-2 secretion most probably enhance the expression of tight junctions in the intestinal wall, with the consequence of decreased gut permeability and leakage of endotoxin (8). It has been suggested that low-grade metabolic inflammation driven by endotoxin translocation is the proposed mechanism by which the microbiota may contribute to systemic inflammation in obesity due to increased intestinal permeability (7,8). *Lactobacilli* may have the capability to improve intestinal integrity in rodents, which may diminish the LPS overflow from the gut to the circulation and

thereby reduce the systemic concentrations of inflammatory markers (42). In our study, LPS and systemic cytokine concentrations remained stable.

Limitations of our study are the small-sized cohorts and the rather short intervention period. However, contrary to other studies in the field, where healthy control subjects and patients with metabolic disorders were analyzed together, we investigated healthy people without diabetes and extensively characterized them metabolically by validated methods including euglycemic-hyperinsulinemic clamp test with isotopic dilution to assess tissue-specific insulin sensitivity and substrate oxidation, different calculations of insulin sensitivity and  $\beta$ -cell function.

Although our study failed to detect physiological consequences of an increased incretin, insulin and C-peptide response in the *L. reuteri* group, these cannot be excluded. With a treatment duration of 4 weeks, our study was not designed to test for long-term effects and changes in body weight were not expected. The slightly increased incretin and insulin concentrations are within the physiological range of metabolic responses, which will not likely affect glucose tolerance of healthy humans. However, it remains to be tested whether patients with type 2 diabetes would respond differently than healthy volunteers.

In conclusion, administration of probiotic *L. reuteri* increased insulin secretion and incretin release in humans. This effect was not caused by a modulation of the fecal microbiota, suggesting a direct effect of the *Lactobacilli* on host physiology. The increase of the intestinotrophic gut peptide, GLP-2, which may contribute to intestinal integrity, was accompanied by stable concentrations of endotoxin and immune mediators. Therefore, administration of a specific bacterial strain might have clinical implications by improving incretin-mediated  $\beta$ -cell function in individuals with impaired glucose homeostasis and therefore warrants further studies on specific bacterial strains in type 2 diabetes.

---

**Acknowledgments.** The authors thank Dr. Achim Huebinger and Dr. Gilles Sequaris for clinical and technical support and the team of the Clinical Research Center of the Institute for Clinical Diabetology, German Diabetes Center.

**Funding.** The study was funded in part by a grant of Heinrich-Heine University and the German Center for Diabetes Research and an unrestricted grant by the Danone Institute for Nutrition and Health e.V., Germany. Nutraceutix provided *L. reuteri* caplets and placebos for this study.

**Duality of Interest.** N.C.S. is employed at Lilly Deutschland, Bad Homburg, Germany. No other potential conflicts of interest relevant to this article were reported.

**Author Contributions.** M.-C.S., H.K., P.S., N.C.S., and M.R. designed the study. M.-C.S., B.N., and F.Z. performed the research. M.-C.S., B.N., P.N., V.B., F.Z., J.-H.H., G.P., B.H., J.J.H., C.M.K., L.B.B., I.M., J.W., and B.H. generated data. M.-C.S., K.S., and I.M. performed the statistical analysis. M.-C.S., H.K., N.C.S., and M.R. drafted the manuscript. All authors critically reviewed the manuscript. M.R. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Prior Presentation.** Parts of this study were presented in abstract form at the 72nd Scientific Sessions of the American Diabetes Association, Philadelphia, PA, 8–12 June 2012, and at The Microbiome (Q8), Keystone, CO, 4–9 March 2012.

## References

- Tabák AG, Jokela M, Akbaraly TN, Brunner EJ, Kivimäki M, Witte DR. Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. *Lancet* 2009;373:2215–2221
- Vilshøj T, Krarup T, Deacon CF, Madsbad S, Holst JJ. Reduced postprandial concentrations of intact biologically active glucagon-like peptide 1 in type 2 diabetic patients. *Diabetes* 2001;50:609–613
- Kolb H, Mandrup-Poulsen T. The global diabetes epidemic as a consequence of lifestyle-induced low-grade inflammation. *Diabetologia* 2010;53:10–20
- Roden M. Mechanisms of Disease: hepatic steatosis in type 2 diabetes—pathogenesis and clinical relevance. *Nat Clin Pract Endocrinol Metab* 2006;2:335–348
- Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 2006;444:1027–1031
- Larsen N, Vogensen FK, van den Berg FWJ, et al. Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. *PLoS ONE* 2010;5:e9085
- Cani PD, Amar J, Iglesias MA, et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 2007;56:1761–1772
- Cani PD, Possemiers S, Van de Wiele T, et al. Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. *Gut* 2009;58:1091–1103
- Million M, Angelakis E, Paul M, Armougoum F, Leibovici L, Raoult D. Comparative meta-analysis of the effect of *Lactobacillus* species on weight gain in humans and animals. *Microb Pathog* 2012;53:100–108
- Silk DB, Davis A, Vulevic J, Tzortzis G, Gibson GR. Clinical trial: the effects of a trans-galactooligosaccharide prebiotic on faecal microbiota and symptoms in irritable bowel syndrome. *Aliment Pharmacol Ther* 2009;29:508–518
- Cani PD, Lecourt E, Dewulf EM, et al. Gut microbiota fermentation of prebiotics increases satietogenic and incretin gut peptide production with consequences for appetite sensation and glucose response after a meal. *Am J Clin Nutr* 2009;90:1236–1243
- Vrieze A, Van Nood E, Holleman F, et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 2012;143:913–916, e7
- Andreasen AS, Larsen N, Pedersen-Skovsgaard T, et al. Effects of *Lactobacillus acidophilus* NCFM on insulin sensitivity and the systemic inflammatory response in human subjects. *Br J Nutr* 2010;104:1831–1838
- Ejtahed HS, Mohtadi-Nia J, Homayouni-Rad A, Niafar M, Asghari-Jafarabadi M, Mofid V. Probiotic yogurt improves antioxidant status in type 2 diabetic patients. *Nutrition* 2012;28:539–543
- Poutahidis T, Kleiweietfeld M, Smillie C, et al. Microbial reprogramming inhibits Western diet-associated obesity. *PLoS ONE* 2013;8:e68596
- Gøbel RJ, Larsen N, Jakobsen M, Mølgaard C, Michaelsen KF. Probiotics to adolescents with obesity: effects on inflammation and metabolic syndrome. *J Pediatr Gastroenterol Nutr* 2012;55:673–678
- Larsen N, Vogensen FK, Gøbel RJ, et al. Effect of *Lactobacillus salivarius* Ls-33 on fecal microbiota in obese adolescents. *Clin Nutr* 2013;32:935–940
- Qin J, Li Y, Cai Z, et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 2012;490:55–60
- Karlsson FH, Tremaroli V, Nookaew I, et al. Gut metagenome in European women with normal, impaired and diabetic glucose control. *Nature* 2013;498:99–103
- Jacobsen CN, Rosenfeldt Nielsen V, Hayford AE, et al. Screening of probiotic activities of forty-seven strains of *Lactobacillus* spp. by in vitro techniques and evaluation of the colonization ability of five selected strains in humans. *Appl Environ Microbiol* 1999;65:4949–4956
- Segal KR, Van Loan M, Fitzgerald PI, Hodgdon JA, Van Itallie TB. Lean body mass estimation by bioelectrical impedance analysis: a four-site cross-validation study. *Am J Clin Nutr* 1988;47:7–14
- Ziegler D, Schadealdt P, Pour Mirza A, et al. [<sup>13</sup>C]octanoic acid breath test for non-invasive assessment of gastric emptying in diabetic patients: validation and relationship to gastric symptoms and cardiovascular autonomic function. *Diabetologia* 1996;39:823–830
- Pacini G, Mari A. Methods for clinical assessment of insulin sensitivity and beta-cell function. *Best Pract Res Clin Endocrinol Metab* 2003;17:305–322
- Mari A, Pacini G, Murphy E, Ludvik B, Nolan JJ. A model-based method for assessing insulin sensitivity from the oral glucose tolerance test. *Diabetes Care* 2001;24:539–548
- Ahrén B, Pacini G. Importance of quantifying insulin secretion in relation to insulin sensitivity to accurately assess beta cell function in clinical studies. *Eur J Endocrinol* 2004;150:97–104
- Kautzky-Willer A, Krssak M, Winzer C, et al. Increased intramyocellular lipid concentration identifies impaired glucose metabolism in women with previous gestational diabetes. *Diabetes* 2003;52:244–251
- Nauack MA, El-Ouaghli A, Gabrys B, et al. Secretion of incretin hormones (GIP and GLP-1) and incretin effect after oral glucose in first-degree relatives of patients with type 2 diabetes. *Regul Pept* 2004;122:209–217
- Nowotny B, Zahiragic L, Krog D, et al. Mechanisms underlying the onset of oral lipid-induced skeletal muscle insulin resistance in humans. *Diabetes* 2013;62:2240–2248
- Krssak M, Falk Petersen K, Dresner A, et al. Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a <sup>1</sup>H NMR spectroscopy study. *Diabetologia* 1999;42:113–116
- Yang S, Lin S, Kelen GD, et al. Quantitative multiprobe PCR assay for simultaneous detection and identification to species level of bacterial pathogens. *J Clin Microbiol* 2002;40:3449–3454
- Byun R, Nadkarni MA, Chhour KL, Martin FE, Jacques NA, Hunter N. Quantitative analysis of diverse *Lactobacillus* species present in advanced dental caries. *J Clin Microbiol* 2004;42:3128–3136
- Haarman M, Knol J. Quantitative real-time PCR analysis of fecal *Lactobacillus* species in infants receiving a prebiotic infant formula. *Appl Environ Microbiol* 2006;72:2359–2365
- Pfleger C, Mortensen HB, Hansen L, et al. Hvidøre Study Group on Childhood Diabetes. Association of IL-1ra and adiponectin with C-peptide and remission in patients with type 1 diabetes. *Diabetes* 2008;57:929–937
- Habich C, Kempe K, van der Zee R, et al. Heat shock protein 60: specific binding of lipopolysaccharide. *J Immunol* 2005;174:1298–1305
- Ristow M, Zarse K, Oberbach A, et al. Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc Natl Acad Sci U S A* 2009;106:8665–8670
- Orskov C, Rabenhøj L, Wettergren A, Kofod H, Holst JJ. Tissue and plasma concentrations of amidated and glycine-extended glucagon-like peptide I in humans. *Diabetes* 1994;43:535–539
- Hartmann B, Johnsen AH, Orskov C, Adelhorst K, Thim L, Holst JJ. Structure, measurement, and secretion of human glucagon-like peptide-2. *Peptides* 2000;21:73–80
- Martínez I, Müller CE, Walter J. Long-term temporal analysis of the human fecal microbiota revealed a stable core of dominant bacterial species. *PLoS ONE* 2013;8:e69621
- Wichmann A, Allahyar A, Greiner TU, et al. Microbial modulation of energy availability in the colon regulates intestinal transit. *Cell Host Microbe* 2013;14:582–590
- Loguercio C, Federico A, Tuccillo C, et al. Beneficial effects of a probiotic VSL#3 on parameters of liver dysfunction in chronic liver diseases. *J Clin Gastroenterol* 2005;39:540–543
- O'Connor LM, Lentjes MA, Luben RN, Khaw KT, Wareham NJ, Forouhi NG. Dietary dairy product intake and incident type 2 diabetes: a prospective study using dietary data from a 7-day food diary. *Diabetologia* 2014;57:909–917
- Mao Y, Nobaek S, Kasravi B, et al. The effects of *Lactobacillus* strains and oat fiber on methotrexate-induced enterocolitis in rats. *Gastroenterology* 1996;111:334–344